

Parent Acknowledgement Letter

Bright Star Academy

3050 Oceanside Road
Oceanside, NY 11572
(516)766-5699

I acknowledge that I have met with Bright Star Academy's Director and /or office staff and my child's teacher regarding his/her healthcare/action plan regarding his/her allergy. I reviewed how to recognize the symptoms and how to administer the medication.

I acknowledge and understand the I am responsible to complete all of the required and mandated forms every 6 months. Failure to do so may result in the immediate dismissal of your child from Bright Star Academy for failure to comply with our plan.

Child's name _____

Parent/Guardian Signature _____

Date _____

Bright Star Academy Staff _____

Bright Star Academy
 3050 Oceanside Road
 Oceanside, NY 11572
 (516) 766-5699

Health Care Plan Packet

Dear Guardian,

All medication forms for your child's allergies and/or special needs must be renewed every six (6) months according to New York State Regulations. Your child's forms must be renewed by _____.

Items that are checked are needed:

Form	Parent signature required	Physician signature required
Form 7006	✓	
Form 6929 (3 pages)	✓	✓
Form 7002 (Medication Consent Form)	✓	✓
Action Plan From Doctor	✓	✓

Form 7002 is required for each medication the child needs. All forms must match the medication given and the dosage. For example, Benadryl, Auvi-Q, and Zyrtec all require their own form.

Medication:

Over The Counter Medication: must be labeled with child's first and last name and have a dosing spoon/cup/dropper that matches the medication amount to be given to the child.

Prescription Medication: must be kept in the original container and have the original pharmacy label on it. Medication will not be accepted without the pharmacy label including child's name, medication name and dosage.

Meals/Snack:

Bright Star Academy will not provide any meals/snack to anyone with a healthcare plan/ allergy/ dietary restriction. All meals, including those at celebrations, must be provided by the family.

Please note: Your child will not be allowed back to school if forms are not updated.

Child's Name _____

Date _____

Date of Plan: / /

THE FOLLOWING STEPS WILL BE TAKEN IF THE CHILD EXHIBITS SYMPTOMS including, but not limited to:

- Inject epinephrine immediately and note the time when the first dose is given.
- Call 911/local rescue squad (Advise 911 the child is in anaphylaxis and may need epinephrine when emergency responders arrive).
- Lay the person flat, raise legs, and keep warm. If breathing is difficult or the child is vomiting, allow them to sit up or lie on their side.
- If symptoms do not improve, or symptoms return, an additional dose of epinephrine can be given in consultation with 911/emergency medical technicians.
- Alert the child's parents/guardians and emergency contacts.
- After the needs of the child and all others in care have been met, immediately notify the office.

MEDICATION/DOSES

- Epinephrine brand or generic:
- Epinephrine dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM

ADMINISTRATION AND SAFETY INFORMATION FOR EPINEPHRINE AUTO-INJECTORS

When administering an epinephrine auto-injector follow these guidelines:

- Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than the mid-outer thigh. If a staff member is accidentally injected, they should seek medical attention at the nearest emergency room.
- If administering an auto-injector to a young child, hold their leg firmly in place before and during injection to prevent injuries.
- Epinephrine can be injected through clothing if needed.
- Call 911 Immediately after injection.

STORAGE OF EPINEPHRINE AUTO-INJECTORS

- All medication will be kept in its original labeled container.
- Medication must be kept in a clean area that is inaccessible to children.
- All staff must have an awareness of where the child's medication is stored.
- Note any medications, such as epinephrine auto-injectors, that may be stored in a different area.
- Explain here where medication will be stored:

MAT/EMAT CERTIFIED PROGRAMS ONLY

Only staff listed in the program's Health Care Plan as medication administrant(s) can administer the following medications. Staff must be at least 18 years old and have first aid and CPR certificates that cover all ages of children in care.

- Antihistamine brand or generic:
- Antihistamine dose:
- Other (e.g., inhaler-bronchodilator if wheezing):

***Note: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.**

STORAGE OF INHALERS, ANTIHISTAMINES, BRONCHODILATOR

All medication will be kept in its original labeled container. Medication must be kept in a clean area that is inaccessible to children. All staff must have an awareness of where the child's medication is stored. Explain where medication will be stored. Note any medications, such as asthma inhalers, that may be stored in a different area.

Explain here:

NEW YORK STATE
OFFICE OF CHILDREN AND FAMILY SERVICES
MEDICATION CONSENT FORM
CHILD DAY CARE PROGRAMS

PARENT COMPLETE THIS SECTION (#19 - #23)

19. If Section #7A is completed, do the instructions indicate a specific time to administer the medication? (For example, did the licensed authorized prescriber write 12pm?) Yes N/A No

Write the specific time(s) the child day care program is to administer the medication (i.e.: 12 pm): _____

20. I, parent, authorize the day care program to administer the medication, as specified on the front of this form, to (child's name): _____

21. Parent's Name (please print): _____	22. Date Authorized: / /
---	--------------------------------

23. Parent's Signature:
X

CHILD DAY CARE PROGRAM COMPLETE THIS SECTION (#24 - #30)

24. Program Name: Bright Star Academy	25. Facility ID Number: 658827	26. Program Telephone Number: (516)766-5699
--	-----------------------------------	--

27. I have verified that (#1 - #23) and if applicable, (#33 - #36) are complete. My signature indicates that all information needed to give this medication has been given to the day care program.

28. Staff's Name (please print): _____	29. Date Received from Parent: / /
--	--

30. Staff Signature:
X

ONLY COMPLETE THIS SECTION (#31 - #32) IF THE PARENT REQUESTS TO DISCONTINUE THE MEDICATION PRIOR TO THE DATE INDICATED IN (#15)

31. I, parent, request that the medication indicated on this consent form be discontinued on _____ / _____ / _____ (Date)

Once the medication has been discontinued, I understand that if my child requires this medication in the future, a new written medication consent form must be completed.

32. Parent Signature:
X

LICENSED AUTHORIZED PRESCRIBER TO COMPLETE, AS NEEDED (#33 - #35)

33. Describe any additional training, procedures or competencies the day care program staff will need to care for this child.

34. Since there may be instances where the pharmacy will not fill a new prescription for changes in a prescription related to dose, time or frequency until the medication from the previous prescription is completely used, please indicate the date you are ordering the change in the administration of the prescription to take place.

DATE: _____ / _____ / _____

By completing this section, the day care program will follow the written instruction on this form and not follow the pharmacy label until the new prescription has been filled.

35. Licensed Authorized Prescriber's Signature:
X



Name: _____ D.O.B.: _____

Allergic to: _____

Weight: _____ lbs. Asthma: Yes (higher risk for a severe reaction) No



NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____

THEREFORE:

- If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:
SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

OR A COMBINATION of symptoms from different body areas.



1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE


PHYSICIAN/HCP AUTHORIZATION SIGNATURE

DATE




HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps; you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPI™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPI by finger grips only and slowly insert the needle into the thigh. SYMJEPI can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____
 DOCTOR: _____ PHONE: _____
 PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____
 NAME/RELATIONSHIP: _____ PHONE: _____
 NAME/RELATIONSHIP: _____ PHONE: _____